

E-Scribe #: 2433693



GaltDirect has partnered with Sterling Specialty Pharmacy **for the easiest and most hassle-free Norgestic® prescribing experience.**

PHONE: 888-618-4126 FAX: 866-588-0371 NPI #: 1225548480
EMAIL: support@sterlingspecialtyrx.com
ADDRESS: 1312 Northland Drive, Suite 500, Mendota Heights, MN 55120

E-Scribe Rx to Sterling Specialty Pharmacy:

STERLING BENEFITS INVESTIGATION:

1. Verify patient's insurance coverage
2. Text or call the patient

IF PATIENT HAS COMMERCIAL COVERAGE:

Copay assistance is applied, patient pays as little as \$0 for their prescription of Norgestic®. Rx is mailed to the patient.*

Rx DENIED OR PATIENT HAS NO COVERAGE:

Rx will be fulfilled by our Patient Assistance Program. Patients will not pay more than \$60 for their Norgestic® prescription. Rx is mailed to the patient.

*Shipping charges may apply.

See the reverse side for important safety information or visit norgestic.com for more details and full prescribing information.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit MedWatch or call 1-800-FDA-1088.

Benefits of Using GaltDirect:



Avoid treatment delays



Increase adherence



Rx mailed to the patient's home (in all 50 states).
Guaranteed delivery within 2 business days.



Reduce the number of phone calls to and from different pharmacies



INDICATIONS AND USAGE

Norgesic (orphenadrine citrate, aspirin and caffeine 25 mg/ 385 mg/ 30 mg)

Tablets are indicated in:

1. Symptomatic relief of mild to moderate pain of acute musculoskeletal disorders.
2. The orphenadrine component is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute painful musculoskeletal conditions.

The mode of action of orphenadrine has not been clearly identified, but may be related to its analgesic properties. Norgesic Tablets do not directly relax tense muscles in man.

IMPORTANT SAFETY INFORMATION (ISI)

Contraindications

Norgesic is contraindicated in patients with glaucoma, pyloric or duodenal obstruction, achalasia, prostatic hypertrophy, obstructions at the bladder neck, myasthenia gravis, hypersensitivity to the drug, and known sensitivities to aspirin or caffeine.

Warnings and Precautions

- Norgesic is not recommended in patients with chicken pox, influenza, or flu symptoms due to the possible development of Reye's Syndrome with aspirin.
- Patients should be cautioned on possible impairment of the ability to engage in potentially hazardous activities such as operating machinery or driving a motor vehicle.
- Aspirin should be used with extreme caution in the presence of peptic ulcers and coagulation.
- Use of NSAIDs, including aspirin, can cause premature closure of the fetal ductus arteriosus and fetal renal dysfunction leading to oligohydramnios and, in some cases, neonatal renal impairment. Because of these risks, limit Norgesic use between about 20 and 30 weeks of gestation, and avoid use at about 30 weeks of gestation and later in pregnancy.
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) has been reported in patients taking NSAIDs such as aspirin. Advise patients to stop taking the

medication immediately if they develop any type of rash or fever and to contact their healthcare provider as soon as possible.

- The safe and effective use in children has not been established; Usage in children under 12 is not recommended.
- Safety of continuous long-term therapy with Norgesic has not been established; If prescribed for prolonged use, periodic monitoring of blood, urine, and liver function values is recommended.
- Confusion, anxiety, and tremors have been reported with concomitant propoxyphene and orphenadrine.

Adverse Reactions

The side effects of Norgesic are those seen with aspirin and caffeine or those usually associated with mild anti-cholinergic agents. These may include tachycardia, palpitation, urinary hesitancy or retention, dry mouth, blurred vision, dilation of the pupil, increased intraocular tension, weakness, nausea, vomiting, headache, dizziness, constipation, and drowsiness, and rarely, urticaria and other dermatosis.